

General Disclaimer


One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

NASW - 3966

DAA/HQ

der



(NASA-CR-176398) EXTENDED ATMOSPHERES OF
OUTER PLANET SATELLITES AND COMETS Interim
Report, Aug. - Oct. 1985 (Atmospheric and
Environmental Research) 10 p HC A02/MF A01

N86-14209

Unclas

CSCL 03B G3/90 04866



ATMOSPHERIC AND ENVIRONMENTAL RESEARCH, INC.
CAMBRIDGE, MASSACHUSETTS

Extended Atmospheres of Outer Planet Satellites
and Comets

✓
William H. Smyth and
Michael R. Combi

Atmospheric and Environmental Research, Inc.
840 Memorial Drive
Cambridge, MA 02139

November 1985

Interim Report for the Period
August 1, 1985 to October 31, 1985

Prepared for
NASA Headquarters

TECHNICAL REPORT STANDARD TITLE PAGE

1. Report No.	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Extended Atmospheres of Outer Planet Satellites and Comets		5. Report Date November 1985	6. Performing Organization Code
		8. Performing Organization Report No.	
7. Author(s) William H. Smyth and Michael R. Combi		10. Work Unit No.	
9. Performing Organization Name and Address Atmospheric and Environmental Research, Inc. 840 Memorial Drive Cambridge, MA 02139		11. Contract or Grant No. NASW-3966	
		13. Type of Report and Period Covered Interim Report August - October 1985	
12. Sponsoring Agency Name and Address NASA Headquarters Headquarters Contract Division Washington, DC 20546		14. Sponsoring Agency Code HW-2	
15. Supplementary Notes			
<p>16. Abstract</p> <p>The new cometary hydrogen particle-trajectory model, completed last year, has been used successfully to analyze observations of Comet P/Giacobini-Zinner. The Pioneer Venus Orbiter Ultraviolet Spectrometer observed the comet at 1216 Å (hydrogen Lyman-α) on 11 September 1985 when the comet was 1.03 AU from the Sun and 1.09 AU from Venus. The analysis implies a production rate at 1.03 AU of $2.3 \times 10^{28} \text{ s}^{-1}$ of the water molecules which photodissociate to produce the observed hydrogen. An upper limit for the H₂O production rate of Comet P/Halley of $5 \times 10^{28} \text{ s}^{-1}$ at 2.60 AU was also obtained from the Pioneer Venus instrument.</p>			
17. Key Words (Selected by Author(s)) satellites comets		18. Distribution Statement	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 8	22. Price*

*For sale by the Clearinghouse for Federal Scientific and Technical Information, Springfield, Virginia 22151.

I. Program of Research for the First Quarter

Research activities during the first quarter have concentrated on the comet portion of this project. Analysis of successful hydrogen observations of Comet P/Giacobini-Zinner by the Pioneer Orbiter Ultraviolet Spectrometer (PVOUS) is discussed below. A paper summarizing these observations and their analysis was presented in October at the AAS/DPS meeting (Stewart, Combi, and Smyth 1985). This represents the second step in our Comet Observing and Analysis Program, a collaborative effort with A.I.F. Stewart, Principal Investigator of PVOUS. The first step involved the successful observations and analysis of Comet P/Encke reported earlier (Combi, Stewart, and Smyth 1984). Nearly continuous observations of Comet P/Halley and their subsequent analysis, to begin in the next quarter, will initiate the third step of this program. From limited observations on September 11, Stewart derived a 3s upper limit of $5 \times 10^{28} \text{ s}^{-1}$ for the H_2O production rate from a null PVOUS observation of Lyman- α in P/Halley when it was 2.60 AU from the Sun. This is consistent with a vectorial model analysis of IUE OH observations on 12.9 September which implied a production rate of $2 \times 10^{28} \text{ s}^{-1}$ (Feldman et al. 1985).

Analysis of Pioneer Venus Observations of Comet P/Giacobini-Zinner

The Pioneer Venus Orbiter Ultraviolet Spectrometer (PVOUS) observed Comet P/Giacobini-Zinner at 1216 / (hydrogen Lyman- α) on 11 September 1985, when the comet was 1.03 AU from the Sun and 1.09 AU from Venus. The field of view of the instrument made 2500 passes across the coma, taking samples every $4.4 \times 10^5 \text{ km}$ along a swath whose width was $1.7 \times 10^6 \text{ km}$. The path of the swath made an angle of about 47° with the comet-sun line as projected on the sky plane. The maximum observed signal was 405 R, including an interplanetary background of 261 R. The fully calibrated and reduced data were provided to us by A.I.F. Stewart.

These data have been analyzed with the AER particle-trajectory model (PTM) based on the production of H atoms by photodissociation of H_2O and OH. This model, developed under this and previous NASA contracts, calculates exact atom trajectories in three dimensions taking into account both variable solar radiation pressure and solar gravity, and includes the fully time-dependent spatially extended random ejection directions of photodissociated atoms and radicals.

The numerical parameters of the model are the same as those used previously for the Lyman- α P/Encke observations (Combi, Stewart, and Smyth 1984) and are scaled in the PTM appropriately for the orbit of P/Giacobini-Zinner. Therefore, all time, heliocentric distance, and heliocentric velocity dependences are automatically taken into account. The H_2O production rate was assumed to vary as the inverse cube of the heliocentric distance, as is appropriate for H_2O sublimation near 1 AU.

Figure 1 shows a comparison of the observed Lyman- α scan with the modeled scan. The model output includes the trapezoidal slit function appropriate to the PVOUS data. The model analysis implies a water production rate of 2.3×10^{28} molecules s^{-1} from Comet P/Giacobini-Zinner at the time of the observation. This value is bracketed by the OH production rate of P/Giacobini-Zinner obtained by 18 cm radio observations by Bockelee-Morvan et al. (1985) in August (1.8×10^{28} molecules s^{-1}) and October (2.9×10^{22} molecules s^{-1}). Contour plots of the model two-dimensional Lyman- α intensity as seen by the Pioneer Venus Orbiter as well as seen normal to the comet plane are shown respectively in Figure 2 and Figure 3. Because of the viewing geometry, the distortion of the hydrogen coma due to solar radiation pressure is more obvious in Figure 3 than in Figure 2.

II. Program of Research for the Second Quarter

Research activities during the second quarter in the area of comets will involve preparing model results to be used for initial interpretation of Comet P/Halley data to be obtained by the Pioneer Venus Spacecraft by A.I.F. Stewart starting in late December. Analysis of the PVOUS data will likely be initiated in January 1986 as it is available from Stewart. Additional effort in the second quarter will also be directed toward writing a paper to document the AER particle-trajectory model.

Research activities in the Saturn system will be continued in the second quarter. Significantly improved values of the electron and ion number densities and temperatures in the Saturn magnetosphere are expected early in the second quarter from E.C. Sittler as part of our continuing collaborative effort with him. This information will be used to refine in our Titan hydrogen torus model the lifetime description of H atoms in the Saturn magnetosphere. Exploratory model computations will then follow as we initiate our analysis of the Lyman- α data acquired by the Voyager 1 and 2 spacecrafts and recently properly formatted by D.E. Shemansky for this purpose.

References

- Bockelee-Morvan, D., Bourgois, G., Crovisier, J., and Gérard, E. (1985) Radio Observations of OH in P/Halley and P/Giacobini-Zinner. Poster 13.18 presented at the AAS/DPS meeting Oct. 28 - Nov. 1, 1985, Baltimore, MD.
- Combi, M.R., Stewart, A.I.F., and Smyth, W.H. (1984) An Analysis of Pioneer Venus Hydrogen Lyman- α Observations of Comet P/Encke. BAAS 16, 638.
- Delsemme, A.H. (1982), Chemical Composition of Cometary Nuclei, in Comets, (L. Wilkening, ed.), University of Arizona Press, Tucson, AZ, p. 85.
- Feldman, P.D., et al. (1985) IUE Observations of Comet P/Halley in September 1985. BAAS 17 (late paper abstract).
- Rottman, G.J. (1985) Private communication.
- Stewart, A.I.F., Combi, M.R., and Smyth, W.H. (1985) Lyman-alpha Measurements of P/Giacobini-Zinner. BAAS 17, 686.





Figure 1

A comparison of the Pioneer Venus observations of Lyman- α from Comet P/Giacobini-Zinner (points) and the PTM (line) analysis is shown. The best model fit yields an H_2O production rate of $2.3 \times 10^{28} \text{ s}^{-1}$ at 1.03 AU and is consistent with a hydrogen atom lifetime of $9.4 \times 10^5 \text{ s}$ at 1 AU.

OBSERVER VIEW INTENSITY (RAYLEIGHs)




DISTANCE FROM THE NUCLEUS (BOX SIZE= 1.25×10^6 KM)

Figure 2

A 2-D map of H Lyman- α from the PTM analysis is shown with the Pioneer Venus Swath. North is down and East is to the left.

TOP VIEW INTENSITY (RAYLEIGHS)



DISTANCE FROM THE NUCLEUS (BOX SIZE= $1.25\text{E}+06\text{KM}$)

Figure 3

An out-of-plane view of the 2-D H Lyman- α map from the PTM analysis is shown. This view maximizes the apparent radiation pressure distortion.

[Handwritten signature]